

## The effects of body size and orientation on ultraviolet radiation exposure

**Link to publication record in USC Research Bank:**

<http://research.usc.edu.au/vital/access/manager/Repository/usc:16469>

**Document Version:**

Author accepted manuscript (postprint)

**Citation for published version:**

Parisi, A V, Kimlin, Michael G, Wong, J C F, Fleming, R A (1996) The effects of body size and orientation on ultraviolet radiation exposure. *Photodermatology Photoimmunology and Photomedicine*, Vol. 12, No. 2, pp.66-72.

**Copyright Statement:**

Copyright © 1996 Wiley-Blackwell Publishing Inc. This is the peer reviewed version of the following article: Parisi, A V, Kimlin, Michael G, Wong, J C F, Fleming, R A (1996) The effects of body size and orientation on ultraviolet radiation exposure. *Photodermatology Photoimmunology and Photomedicine*, Vol. 12, No. 2, pp.66-72, which has been published in final form at <http://dx.doi.org/10.1111/j.1600-0781.1996.tb00177.x>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

**General Rights:**

Copyright for the publications made accessible via the USC Research Bank is retained by the author(s) and / or the copyright owners and it is a condition of accessing these publications that users recognize and abide by the legal requirements associated with these rights.

**Take down policy**

The University of the Sunshine Coast has made every reasonable effort to ensure that USC Research Bank content complies with copyright legislation. If you believe that the public display of this file breaches copyright please contact [research-repository@usc.edu.au](mailto:research-repository@usc.edu.au) providing details, and we will remove the work immediately and investigate your claim.

# THE EFFECTS OF BODY SIZE AND ORIENTATION ON ULTRAVIOLET RADIATION EXPOSURE

A.V. Parisi<sup>\*</sup>, M.G. Kimlin<sup>\*</sup>, J.C.F. Wong<sup>†</sup> and R.A. Fleming<sup>†</sup>

<sup>†</sup>School of Physics, Queensland University of Technology, GPO Box 2434,  
BRISBANE 4001 AUSTRALIA

<sup>\*</sup>Centre for Astronomy and Atmospheric Research, Faculty of Sciences, University of  
Southern Queensland, TOOWOOMBA 4350 AUSTRALIA

**Abstract**

A method has been developed for determining the UV and erythematous exposures to the entire body. The difference between the ambient erythematous exposure and that to the body compared to the ambient exposure may be as high as 76%. The height, orientation, and overall height had a minimal effect on the exposure to the body with size, time of day and time of year having a significant effect. The diffuse component of UV to a side of the body ranged from 20% to 41% between different times of the year with different levels of cloud cover. The ratio of the body to the ambient erythematous exposures varied from 0.24 to 0.61 with the time of day and time of year with the smaller value for periods of high solar altitude.

**Keywords:** UV dosimeters; polysulphone; erythema; effects on exposure; exposures to the body

## Introduction

Decreased levels of stratospheric ozone (1) and increased levels of terrestrial UVB (2,3) have been measured. These increased levels of UVB have generated concerns about the resultant deleterious effects on human health. Reviews in the literature list the UV induced immediate reactions as erythema, photodermatoses, keratitis and conjunctivitis with the longer term effects of skin cancer, photo-aging of the skin and cataracts along with the damaging effects to the immune system and DNA (4).

Research into the UV induced effects on humans requires measurement of personal UV exposure. A number of studies have employed UV dosimeters to investigate the UV radiation exposures to selected sites on rotating headform models and on human subjects undertaking a variety of activities (5-8). Alternatively, a UV sensor worn on the lapel or waistband and connected to a portable data logger has been employed to monitor UV exposure rate during a number of outdoor activities (9). The effect of the inclination of the receiving plane on the biologically effective UV has also been investigated (10). These studies have provided the UV exposures to specific sites. However, no information has been provided on the exposure to the entire body. Additionally, the exposures with the headform models make the important assumption that there is no effect due to different body size and orientation. This paper presents a method developed to determine the UV exposure to the entire body and to investigate the effects of the body size and orientation on the UV exposure.

The topography, orientations and movements of the human body are complex and practically impossible to model. As a result, a simplified model of the human body

will be employed to study the effects of size and orientation. Polysulphone dosimeters will be deployed at a number of sites on rectangular prisms of various sizes and orientations and which represent models of human bodies. The UV and erythral exposures to each site will be interpolated between sites and summed over the shape to provide the total exposure to the body.

## **Materials and Methods**

### *Calibrations*

Polysulphone in thin film form of the order of 40  $\mu\text{m}$  thick was employed for the UV dosimeters (11-13). The polysulphone dosimeters (supplied by A Davis, 3 Cumley Rd., Toothill, Ongar, Essex, CM5 9SJ, UK) consisted of the film mounted in cardboard holders 30 x 30 mm with an active area of 16 x 12 mm. As a result of UV exposure, the polysulphone photodegrades and this is quantified by measurement of the change in optical absorbance ( $\Delta A$ ) at 330 nm. In this research, to standardise the read out times, the optical absorbance was measured with the spectrophotometer (Shimadzu Co., Kyoto, Japan) immediately pre and post exposure in order to eliminate errors due to a continued increase of absorbance in the dark post exposure (12). For consistency, all of the exposures were performed with the polysulphone on a white backing.

To relate the  $\Delta A$  to the UV exposure, a calibration curve for the polysulphone was obtained by simultaneously exposing the polysulphone and measuring the spectral irradiance with a spectroradiometer in 1 nm intervals on a horizontal unshaded location on a cloud free autumn day. The response of polysulphone does not exactly match the erythral action spectrum, however, the polysulphone may be employed to

measure UV exposures with acceptable accuracy provided they are calibrated. In this paper, the polysulphone dosimeters will be employed to measure autumn solar exposures and the dosimeters have been calibrated against a spectroradiometer to autumn sunshine. The spectroradiometer is based on a double UV holographic grating monochromator (Jobin-Yvon, model DH10, 16-18 rue du canal 91163 France) with calibration traceable to the primary Australian standard lamp housed at the National Measurement Laboratory, CSIRO, Lindfield (14). A number of polysulphone dosimeters were exposed on a horizontal plane within 30 cm of the input aperture of the spectroradiometer from 08:48 to 12:36 Eastern Standard Time (EST). The UV spectrum was measured with the spectroradiometer at 12 minute intervals and a polysulphone dosimeter was removed at each 12 minute interval. As a result, polysulphone dosimeters were exposed for various periods ranging from 12 minutes to 3.8 hours from early morning to noon. The spectral irradiance,  $S(\lambda,t)$  was converted to a UV exposure over a time interval,  $T$ , by:

$$UV = \int_0^T \int_{uv} S(\lambda,t) d\lambda dt \quad (1)$$

where the integration is the summation between the wavelengths 280 to 340 nm. These limits were employed as the solar irradiance is zero at 280 nm and the response of polysulphone is zero at 340 nm (13). The erythema biologically effective UV (UVBE) was also calculated from the spectral irradiance by:

$$UVBE = \int_0^T \int_{uv} S(\lambda,t)A(\lambda)d\lambda dt \quad (2)$$

where  $A(\lambda)$  is the erythema action spectrum (15). Employing Equation (1) provided a calibration for the unweighted UV irradiance from 280 to 340 nm. Equation (2) provided a second calibration curve for the erythema exposures.

In order to compare the exposures of the models of the bodies with the ambient exposures, the output of an IL1400 (International Light, Newburyport, MA) radiometer with a SEL400 photodetector and filter was calibrated against the irradiance measured with the spectroradiometer. The two measurements were made on the same plane and within 30 cm of one another.

### *Preliminary Exposure*

A model of the human body in the shape of a rectangular prism on a stand was exposed to autumn solar radiation in Toowoomba (27.5° S latitude), Australia on 1 March, 1995 for 1 hour between 11:00 and 12:00 EST with an average solar zenith angle of 22°. This was designed as a preliminary exposure to determine the number of dosimeters required on each side. The size of the model was 200 mm x 400 mm with a height of 450 mm. The shape was placed on a stand to provide an overall height of 1350 mm and orientated with the small side facing north. The amount of cloud cover was 3 octas, however no cloud covered the solar disc. Four dosimeters were placed on the top of the shape, nine on each of the larger sides and six on each of the smaller sides. Following exposure, the variation in exposure to each dosimeter on the sides and the top were of the same magnitude as the error, and as a result it is possible to reduce the number of dosimeters over the body to four on each side, with a dosimeter at each corner and one dosimeter on the top side located in the centre. A diagram of the model with the dosimeters attached is provided in Figure 1 with the azimuth angles relative to north, and the inclination angles of the dosimeters on each side in Table 1.

### *Exposures*

Models of the human body with the sizes and orientations in Table 2 were exposed in an open area. To prevent shading, the shapes were spaced several metres apart as shown in Figure 2. The sizes and orientations were selected to investigate the effects of: the height of the body off the ground; the overall height of the body; the shape of the body and the orientation of the body relative to north. The exposures were performed on a cloud free autumn day on 18 April in Toowoomba. The times of the exposures were 09:00 to 10:00, 12:00 to 13:00 and 15:00 to 16:00 EST for the morning, noon and afternoon exposures with average solar zenith angles for each period of  $52^\circ$ ,  $40^\circ$  and  $65^\circ$  respectively.

The exposures at each dosimeter site were employed to calculate the exposure to the entire body (16). The computer software developed divides each side into elements and the exposures at each dosimeter site are interpolated, firstly, vertically and secondly, horizontally to provide the exposure at each element followed by summation to calculate the total exposure to each side. The number of elements in the vertical and horizontal directions was taken as thirty. From previous research (16), this was considered as adequate with a higher number of elements not producing a significant change in the result. For the top side, the exposure at the single site multiplied by the surface area provided the exposure to that side. Summation of the exposures to each side and the top and divided by the total surface area provided the exposure (total exposure/surface area) to the body.

For each period, the ambient irradiance was measured with the calibrated IL1400 radiometer at regular intervals. The IL1400 radiometer was also employed to

determine the UV reflectivity of the ground by measuring the downward and upward fluxes.

## **Results**

### *Calibrations*

The calibration curve of polysulphone for erythema exposure is provided in Figure 3. A similar curve was obtained for calibration for UV exposures between 280 and 340 nm. The slope of these dose response curves decreases for  $\Delta A$  values above 0.3. For values of  $\Delta A$  less than 0.3, previous research in the literature (17) has determined an error of about 10% in the UV exposures obtained.

### *Preliminary Exposure*

The UV and erythema exposures to the body calculated employing the individual exposures at each site are shown Table 3. These exposures are divided by the surface area. The errors in these exposures and in the next Section are taken as 10%. The ambient exposures have been taken as those measured with the dosimeters at the top site of the body. The ratios of the UV and erythema exposures to the body compared to the respective ambient exposures are also shown in Table 3. From these results, the difference between the ambient erythema exposure and the exposure to the body compared to the ambient exposure is 76%.

### *Exposures*

The erythema exposures to each of the sides of the body with the small side facing north for the 1 March between 11:00 and 12:00 EST and the 9.00 to 10:00 and 12:00 to 13:00 EST exposures on 18 April are shown in Table 4. These are the total

exposures to each side calculated from the exposures to the individual sites and divided by the surface area of the respective side. For the 1 March and noon on 18 April, the north side of the body is exposed to sun and the south side is in full shadow. Nevertheless, the south side receives an appreciable amount of erythemal exposure as a result of the high component of diffuse or scattered UVB. On the 1 March, the south side receives 41% compared to the north side and for noon on 18 April, the south side receives 20% compared to the north side. For the morning on 18 April, the east side is in direct sun and the west side is shadowed. The west side receives 23% of the erythemal exposure compared to the east side. The higher diffuse component on 1 March may be attributed to the additional scattering by the three octas cloud cover compared to zero cloud cover on 18 April.

The UV and erythemal exposures to the body for each of the shapes are presented in Table 5 with the erythemal exposures plotted in Figure 4. Shape 3 (square shape) receives a higher exposure in the morning and noon exposures with no significant difference in the afternoon. The larger exposure for the first two periods are most likely due to the square shape providing a larger proportion of its surface area to the direct solar UV. The difference in exposure to this shape is not significant in the afternoon. This is likely to be due to the low solar zenith angle and corresponding low solar irradiances at this time.

The exposures to shapes 2 and 4 show that for the ground cover in this study (dry grass) there is no significant effect due to height of the body and overall height at any time of the day. This is supported by the low measured UV reflectivity off the ground of 2%. There may have been an effect due to height for a ground cover with a higher

ground reflectance (10). The orientation of the shapes had negligible effect on the exposure to the body with only a minimal difference in the exposures to shapes 1, 5 and 6 with the large side facing north, east or north-east.

The ambient UV and UVBE exposures at each period were taken as the average measured with the polysulphone dosimeters at the top side of each body and are provided in Table 6. This Table also provides the ambient UV exposure measured with the IL1400 radiometer and these agree with the dosimeter values within the 10% error margin. For the noon period, the standard error in the ambient UV exposures measured with the dosimeters is 6%. This verifies that the ambient exposure is uniform over the exposure area.

Compared to the noon erythemal ambient exposure, the corresponding morning exposure is less by 54% and the afternoon one is less by 86%. Employing the data in Table 5, the erythemal exposures to the whole body have been averaged over all six shapes for each of the three periods. In contrast to the results for the ambient exposures, compared to the noon erythemal body exposure, the morning exposure is 31% less and the afternoon one is 76% less. From this result, the exposure to the body in the morning is much higher than would have been assumed by measurement of the ambient exposure. A similar result was noted for the UV exposures between 280 and 340 nm. This is due predominantly to the zenith angle of the sun in the morning providing a higher proportion of the exposure to the sides relative to the top. To a lesser extent, other factors that may influence this are variations during the day in the diffuse component of UV due to changes in the transmission properties of the

atmosphere along with any possible changes in UV reflective structures and ground cover.

This variation in the difference between the ambient and the body exposures occurs not only with different times of the day, but also with different times of the year. For example, from Table 3 and Table 5, the erythemal exposure to the body for shape 5 with the small side facing north varies from 26.9 to 19.2 mJ cm<sup>-2</sup> for 1 March and 18 April respectively. In contrast, from Table 4, the ambient erythemal UV varies from 110 to 61 mJ cm<sup>-2</sup>. The ambient UVBE exposure for the one hour period decreased by 45% compared to the 1 March from early to late autumn whereas the UVBE exposure to the body decreased by only 29%.

The ratios of the erythemal exposures to the body compared to the ambient erythemal exposure for the 18 April are shown in Table 7. This ratio changes throughout the day and is significantly lower for every shape for the noon exposure. For example, for shape 5 with the small side facing north, the ratio of the body to the ambient exposure varies from 0.48 to 0.30 to 0.41 for the morning, noon and afternoon exposures respectively. This ratio is also dependent of the time of year as seen by comparing with the data in Table 3 where the value is 0.24 for the exposure between 11:00 and 12:00 on 1 March with the same sized shape and orientation. This even smaller value of the ratio is due to the smaller zenith angle in early autumn compared to the larger angle in late autumn for the other exposures.

## Conclusions

This paper has presented a method developed to determine the UV and erythematous exposures to the entire body. The UV and erythematous exposures were measured with polysulphone dosimeters at a total of 17 sites over a human body model and from these the total exposures to the body have been calculated. The accuracy of these exposures to the body is 10% or better. The difference between the ambient erythematous exposure and the exposure to the body compared to the ambient exposure may be as high as 76%.

The effects of body size and orientation on the UV exposures have been investigated by undertaking a series of exposures of model shapes and determining the UV exposures to the entire body. Measurement of the exposures to the individual sides with different azimuth angles relative to north found a variation of 20% to 41% between different days in the diffuse component of UV. This highlights the complexity and randomness of the incoming UV radiation.

Significant differences occur in the UV and erythematous exposures to the body as a result of the size of the shape, time of the day and the time of the year. The effects of orientation, height and overall height were insignificant on the exposures to the body. The erythematous exposure to a body of the same size and orientation decreased by 29% from early to late autumn whereas the ambient exposure decreased by 45%. Similarly, the erythematous exposure to the body in the morning was 31% less compared to that for the noon period with the ambient exposure 54% less than that at noon. The exposures to the body at times of higher zenith angle are higher than would be assumed by measurement of the ambient exposure.

The ratio of the body to the ambient exposures varied with the time of day and time of year. For the exposures in this research, this ratio varied from 0.24 to 0.61 with the smaller value for periods of high solar altitude, for example, noon. This variation is due to the multifactorial influence of changes in solar zenith and azimuth angles, clouds, transmission properties of the atmosphere and reflective structures and ground cover. This research highlights the differences that exist between the ambient exposures and the exposures to the body and that it is impossible to undertake one measurement with a radiometer or dosimeter at one site and relate this to the exposure to the body.

### **Acknowledgements**

The authors thank Ron Matthews in the mechanical workshop of the Faculty of Sciences, USQ for his assistance.

## References

1. Stolarski R, Bojkov R, Bishop L, Zerefos C, Staehelin J, Zawodny J. Measured trends in stratospheric ozone. *Science* 1992; **256**: 342-349.
2. Blumthaler M, Ambach W. Indication of increasing solar ultraviolet-B radiation flux in alpine regions. *Science* 1990; **248**: 206-208.
3. Kerr J B, McElroy C T. Evidence for large upward trends of ultraviolet-B radiation linked to ozone depletion. *Science* 1993; **262**: 1032-1034.
4. Ambach W, Blumthaler M. Biological effectiveness of solar UV radiation in humans. *Experientia* 1993; **49**: 747-753.
5. Airey D K, Wong J C F, Fleming R A. A comparison of human- and headform-based measurements of solar ultraviolet B dose. *Photodermatol Photoimmunol Photomed* 1995; **11**: 155-158.
6. Herlihy E, Gies P H, Roy C R, Jones M. Personal dosimetry of solar UV radiation for different outdoor activities. *Photochem Photobiol* 1994; **60**: 288-294.
7. Wong C F, Fleming R A, Carter S J, Ring I T, Vishvakarman D. Measurement of human exposure to ultraviolet-B solar radiation using a CR-39 dosimeter. *Health Phys* 1992; **63**: 457-461.
8. Holman C D J, Gibson I M, Stephenson M, Armstrong B K. Ultraviolet irradiation of human body sites in relation to occupation and outdoor activity: field studies using personal UVR dosimeters. *Clin Exp Dermatol* 1983; **8**: 269-277.

9. Diffey B L. Behaviour outdoors and its effects on personal ultraviolet exposure rate measured using an ambulatory datalogging dosimeter. *Photochem Photobiol* 1995: **61**: 615-618.
10. Schaubberger G. Model for the global irradiance of the solar biologically-effective ultraviolet-radiation on inclined surfaces. *Photochem Photobiol* 1990: **52**: 1029-1032.
11. Davis A, Deane G H W, Diffey B L. Possible dosimeter for ultraviolet radiation. *Nature* 1976: **261**: 169-170.
12. Diffey B L. Ultraviolet radiation dosimetry with polysulphone film. In: Diffey B L, ed. *Radiation measurement in photobiology*. New York: Academic Press, 1989: 136-159.
13. CIE (International Commission on Illumination) Technical Report: Personal dosimetry of UV radiation, 1992: Publication No. CIE **98**.
14. Wong C F, Toomey S, Fleming R A, Thomas B W. UV-B radiometry and dosimetry for solar measurements. *Health Phys* 1995: **68**: 175-184.
15. CIE (International Commission on Illumination) Research Note: A reference action spectrum for ultraviolet induced erythema in human skin, 1987: *CIE J* **6**: 17-22.
16. Parisi A V, Wong C F. A dosimetric technique for the measurement of ultraviolet radiation exposure to plants. *Photochem Photobiol* 1994: **60**: 470-474.
17. Diffey B L. A comparison of dosimeters used for solar ultraviolet radiometry. *Photochem Photobiol* 1987: **46**: 55-60.

Table 1 - The dosimeter orientations on each side of the body with the azimuth angles relative to north, and the inclination angles relative to the horizontal.

Side	Dosimeter Orientations	
	Azimuth	Inclination
Top	-	0
North	0	90
East	90	90
South	180	90
West	270	90

Table 2 - The lengths, widths, heights, overall heights and orientations of the six shapes.

Shape Number	Length x width x height (m)	Overall Height (m)	Orientation
1	0.2 x 0.4 x 0.45	1.35	Large side facing North
2	0.2 x 0.4 x 0.45	0.685	Small side facing North
3	0.4 x 0.4 x 0.45	1.35	Side facing North
4	0.2 x 0.4 x 1.20	1.35	Small side facing North
5	0.2 x 0.4 x 0.45	1.35	Small side facing North
6	0.2 x 0.4 x 0.45	1.35	Large side facing NE & SW, smaller side facing NW & SE

Table 3 - The UV and erythemal exposures to the entire body along with the ratio of the body to ambient exposures for the 11:00 to 12:00 EST exposure on 1 March.

UV (J cm <sup>-2</sup> )	UVBE (mJ cm <sup>-2</sup> )	Body/Ambient	
		UV	UVBE
2.37	26.9	0.27	0.24

Table 4 - Erythematous exposures to each of the sides of the body with the small side facing north for the 1 March and the morning and noon exposures on 18 April.

Side	Erythematous Exposure ( $\text{mJ cm}^{-2}$ )		
	1 March (noon)	18 April (morning)	18 April (noon)
Top	110	27	61
North	27	12	35
East	14	22	9
South	11	5	7
West	12	5	9

Table 5 - The UV and erythemat (UVBE) exposures to the entire body for each shape for the morning, noon and afternoon exposures on 18 April.

Shape	Morning		Noon		Afternoon	
	UV	UVBE	UV	UVBE	UV	UVBE
	(J cm <sup>-2</sup> )	(mJ cm <sup>-2</sup> )	(J cm <sup>-2</sup> )	(mJ cm <sup>-2</sup> )	(J cm <sup>-2</sup> )	(mJ cm <sup>-2</sup> )
1	1.17	12.4	1.99	22.2	0.49	4.8
2	1.25	13.2	1.78	19.7	0.57	5.7
3	1.62	17.6	2.08	23.3	0.50	4.9
4	1.36	14.4	1.53	16.7	0.48	4.7
5	1.30	13.8	1.74	19.2	0.40	3.8
6	1.36	14.5	2.03	22.6	0.57	5.6

Table 6 - The erythemal and ambient UV exposures measured with the polysulphone dosimeters at the top side of the bodies and with the calibrated IL1400 radiometer for each of the periods.

Morning			Noon			Afternoon		
Polysulphone		IL1400	Polysulphone		IL1400	Polysulphone		IL1400
UVBE	UV	UV	UVBE	UV	UV	UVBE	UV	UV
(J cm <sup>-2</sup> )								
0.029	2.6	2.6	0.063	5.3	5.0	0.009	0.91	1.0

Table 7 - Ratios of the erythemal exposures to the body compared to the ambient erythemal exposure for the 18 April.

Shape	Body/Ambient		
	Morning	Noon	Afternoon
1	0.43	0.35	0.52
2	0.46	0.31	0.61
3	0.61	0.37	0.53
4	0.50	0.27	0.51
5	0.48	0.30	0.41
6	0.50	0.36	0.60

Figure 1 - Diagram of the model of a body with the dosimeters attached at the selected sites.

Figure 2 - Photograph of the models with the dosimeters.

Figure 3 - Calibration of the polysulphone dosimeters relating the change in absorbance,  $\Delta A$  to the erythematous exposure.

Figure 4 - Erythematous exposures to each body for the (1) morning, (2) noon and (3) afternoon exposures.







