


Graham Smith,^{a*} Urs D.
Wermuth,^b David J. Young^b and
Jonathan M. White^c^aSchool of Physical and Chemical Sciences,
Queensland University of Technology, GPO
Box 2434, Brisbane 4001, Australia, ^bSchool
of Science, Griffith University, Nathan 4111,
Australia, and ^cSchool of Chemistry, University
of Melbourne, Parkville 3052, Australia

Correspondence e-mail: g.smith@qut.edu.au

Key indicators

Single-crystal X-ray study
 $T = 130$ K
Mean $\sigma(\text{C}-\text{C}) = 0.009$ Å
 R factor = 0.063
 wR factor = 0.111
Data-to-parameter ratio = 6.4For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.The β -form of *trans*-3-(2-nitrophenyl)prop-
2-enoic acid at 130 KThe crystal structure of a non-centrosymmetric polymorphic
modification of the title compound, $\text{C}_9\text{H}_7\text{NO}_4$, determined at
130 K, shows the presence of 2_1 screw-generated one-
dimensional zigzag chains formed through catemeric $C(4)$
syn-anti carboxylic acid hydrogen-bonding associations.Received 8 April 2006
Accepted 18 April 2006

Comment

A large number of the *trans*-cinnamic acids, including those
with aromatic ring substituents, undergo topotactic photo-
dimerization reactions (Cohen & Schmidt, 1964; Cohen *et al.*,
1964; Atkinson *et al.*, 2002). The stable α -forms of these acids
give α -truxillic acids, the metastable β -forms give β -truxinic
acids, while the γ -forms give no reaction (Cohen *et al.*, 1964).
Thus irradiation of the title compound, (I) (common name 2-
nitro-*trans*-cinnamic acid), gives the substituted β -truxinic
acid [3,4-bis(2-nitrophenyl)cyclobutane-1,2-dicarboxylic acid].
The unit-cell data for β -2-nitro-*trans*-cinnamic acid were
reported by Schmidt (1964) among a set of 15 β -*trans*-
cinnamic acids, all having one cell dimension between 3.8 and
4.1 Å. This molecular separation allowed their classification as
 β -acids among the three forms, correlating with the two-
dimensional packing which gives the stated 3.6–4.1 Å ethyl-
enic $\text{C}=\text{C}$ separations required for photodimerization to
occur. This reaction proceeds *via* either head-to-tail inter-
action (α -forms) or head-to-head interaction (β -forms)
(Atkinson *et al.*, 2002). Understandably, very few crystal
structures of these acids, including (I), have been previously
reported. Examples of β -acid structures include 4-chloro-
trans-cinnamic acid (Glusker *et al.*, 1975), 4-formyl-*trans*-
cinnamic acid (Nakanishi *et al.*, 1985) and 3-nitro-*trans*-
cinnamic acid (Udaya Lakshmi *et al.*, 2005).We have now determined the structure of (I) using
diffraction data obtained at 130 K from a CCD-detector-
equipped diffractometer and it is reported here. The unit-cell
dimensions and space group determined for (I) are different
from those of Schmidt (1964) for the β form ($a = 3.76$ Å, $b =$
 7.2 Å, $c = 31.5$ Å, $\beta = 91.6^\circ$, space group $P2_1/c$, $Z = 4$). Acid (I)
has comparable cell parameters (but with *ca* $1/2c$ for $Z = 2$)
and may be assumed to represent a non-centrosymmetric




Figure 1
Molecular conformation and atom-naming scheme for (I). Atoms are shown as 40% probability displacement ellipsoids.




Figure 2
Syn-anti hydrogen-bonding associations (broken lines), which extend the structure of (I) into one-dimensional zigzag chains along the *b* cell direction.

polymorph of the room-temperature centrosymmetric β -form of Schmidt. The structure determination shows that the molecules are not planar (Fig. 1), with the nitro group and the essentially planar propenoic acid substituent group both rotated significantly out of the plane of the benzene ring [torsion angles C1–C2–N2–O22 = -154.0 (6°) and C2–C1–C7–C8 = -147.8 (8°)]. However, the molecular conformation is influenced by the presence of intramolecular interactions between the C7 ethylenic H and both a nitro O [C7 \cdots O21 = 2.721 (8) Å and C7–H7 \cdots O21 = 101°] and a carboxyl O [C7 \cdots O91 = 2.767 (8) Å and C7–H7 \cdots O91 = 100°]. The carboxylic acid groups of the 2_1 screw-related molecules associate through single hydrogen bonds [O91 \cdots O92 i = 2.632 (7) Å and O91–H91 \cdots O92 i = 180° ; symmetry code (i) $-x + 2, y - \frac{1}{2}, -z + 1$], giving a *syn-anti* C(4)

catemeric motif (Leiserowitz, 1976) which extends the structure into one-dimensional zigzag chains along the *b* cell direction (Fig. 2). This type of association is found in only a small number of carboxylic acids where, among the mono- and dicarboxylic acids without other functional groups, occurrence of the cyclic $R_2^2(8)$ motif has a probability of 95.5% (Allen *et al.*, 1998). This includes other known photodimerizable β -examples (Glusker *et al.*, 1975; Nakanishi *et al.*, 1985; Udaya Lakshmi *et al.*, 2005). The stacking of the molecules in the chains in (I) provides head-to-head interaction down the short *a*-axial direction, which is consistent with the requirement for β -photodimerization.

Experimental

The title compound was obtained by slow evaporation of a solution of the commercial acid from 80% 2-propanol–water as pale brown crystals [literature m.p. 513 K (Rappoport, 1967)].

Crystal data

$C_9H_7NO_4$	$Z = 2$
$M_r = 193.16$	$D_x = 1.537 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
$a = 3.6850$ (18) Å	$\mu = 0.12 \text{ mm}^{-1}$
$b = 7.074$ (4) Å	$T = 130$ (2) K
$c = 16.073$ (8) Å	Plate, pale brown
$\beta = 94.978$ (11°)	$0.30 \times 0.20 \times 0.03 \text{ mm}$
$V = 417.4$ (4) Å 3	

Data collection

Bruker SMART CCD area-detector diffractometer	810 independent reflections
φ and ω scans	578 reflections with $I > 2\sigma(I)$
Absorption correction: none	$R_{\text{int}} = 0.096$
2152 measured reflections	$\theta_{\text{max}} = 25.0^\circ$

Refinement

Refinement on F^2	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.063$	$w = 1/[\sigma^2(F_o^2) + (0.0161P)^2]$
$wR(F^2) = 0.111$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.81$	$(\Delta/\sigma)_{\text{max}} < 0.001$
810 reflections	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
127 parameters	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$

The carboxylic acid proton was located by difference methods and constrained in the refinement (O–H = 0.90 Å). Other H atoms were included in the refinement at calculated positions (C–H = 0.95 Å) and treated as riding. $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{O})$.

Data collection: SMART (Bruker, 2000); cell refinement: SMART; data reduction: SAINT (Bruker, 1999); program(s) used to solve structure: SHELXS97 (Sheldrick, 1997) in WinGX (Farrugia, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997) in WinGX; molecular graphics: PLATON (Spek, 2003); software used to prepare material for publication: PLATON.

The authors acknowledge financial support from The University of Melbourne, the School of Science, Griffith University, and the School of Physical and Chemical Sciences of the Queensland University of Technology.

References

- Allen, F. H., Raithby, P. R., Shields, G. P. & Taylor, R. (1998). *Chem. Commun.* pp. 1043–1044.

- Atkinson, S. D. M., Almond, M. J., Bowmaker, G. A., Drew, M. G. B., Feltham, E. J., Hollis, P., Jenkins, S. L. & Wiltshire, K. S. (2002). *J. Chem. Soc. Perkin Trans. 2*, pp. 1533–1537.
- Bruker (1999). *SAINT*. Version 6.02. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2000). *SMART*. Version 5.55. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cohen, M. D. & Schmidt, G. M. J. (1964). *J. Chem. Soc.* pp. 1996–2000.
- Cohen, M. D., Schmidt, G. M. J. & Sonntag, F. I. (1964). *J. Chem. Soc.* pp. 2000–2013.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Glusker, J. P., Zacharias, D. E. & Carrel, H. L. (1975). *J. Chem. Soc. Perkin Trans. 2*, pp. 68–74.
- Leiserowitz, L. (1976). *Acta Cryst.* **B32**, 775–802.
- Nakanishi, H., Hasegawa, M. & Mori, T. (1985). *Acta Cryst.* **C41**, 70–71.
- Rappoport, Z. (1967). *Handbook of Tables for Organic Compound Identification*, p. 208. Cleveland, USA: CRC Press Inc.
- Schmidt, G. M. J. (1964). *J. Chem. Soc.* pp. 2014–2021.
- Sheldrick, G. M. (1997). *SHELXL97* and *SHELXS97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Udaya Lakshmi, K., Thamocharan, S., Srinivasan, M., Ramamurthi, K. & Varghese, B. (2005). *Acta Cryst.* **E61**, o3636–o3638.