



---

# Vitamin D insufficiency in south-east Queensland

McGrath, J J; Kimlin, Michael G; Saha, S; et.al.

<https://research.usc.edu.au/esploro/outputs/letter/Vitamin-D-insufficiency-in-south-east-Queensland/99449564002621/filesAndLinks?index=0>

---

McGrath, J. J., Kimlin, M. G., Saha, S., Eyles, D. W., & Parisi, A. V. (2001). Vitamin D insufficiency in south-east Queensland. In *Medical Journal of Australia* (Vol. 174, Issue 3, pp. 150–151). Australasian Medical Publishing Company Pty. Ltd. <https://doi.org/10.5694/j.1326-5377.2001.tb143195.x>  
Document Type: Accepted Version

---

Link to Published Version: <https://doi.org/10.5694/j.1326-5377.2001.tb143195.x>

UniSC Research Bank: <https://research.usc.edu.au>  
[research-repository@usc.edu.au](mailto:research-repository@usc.edu.au)

McGrath JJ, Kimlin MG, Saha S, et al. Vitamin D insufficiency in south-east Queensland. *Med J Aust* 2001; 174(3): 150-151. © Copyright 2001. The Medical Journal of Australia - reproduced with permission.  
Downloaded On 2024/05/16 12:41:57 +1000

*McGrath JJ, Kimlin M, Saha S, Eyles D, Parisi A. (2001) Vitamin D insufficiency in south-east Queensland. Medical Journal of Australia. 174; 150-151.*

## **Vitamin D insufficiency in south-east Queensland**

Associate Professor John J. McGrath MBBS, PhD  
Director, Queensland Centre for Schizophrenia Research  
And Department of Psychiatry, University of Queensland

Mr. Michael G. Kimlin BAppSc, MAppSc  
Associate Lecturer, Physics  
Centre for Astronomy and Atmospheric Research  
Department of Biological and Physical Sciences  
Faculty of Sciences  
University of Southern Queensland

Mr. Sukanta Saha MSc, MCN  
Scientist – Epidemiology Stream  
Queensland Centre for Schizophrenia Research

Dr. Darryl W. Eyles BSc, PhD  
Scientist – Biomedical Stream  
Queensland Centre for Schizophrenia Research

Dr. Alfio V. Parisi PhD, MAppSc  
Senior Lecturer  
Centre for Astronomy and Atmospheric Research  
Department of Biological and Physical Sciences  
Faculty of Sciences  
University of Southern Queensland

Address for correspondence:  
Associate Professor John McGrath  
Queensland Centre for Schizophrenia Research  
Wolston Park Hospital  
Wacol, Queensland, 4076  
Australia  
Ph +61 7 3271 8595  
Fax +61 7 3271 8567  
E-mail [jjm@brain.wph.uq.edu.au](mailto:jjm@brain.wph.uq.edu.au)

Studies from Perth<sup>1</sup>, Hobart<sup>2</sup>, central Victoria<sup>3</sup>, and Adelaide<sup>4</sup> have recently drawn attention to the surprisingly high rates of vitamin D insufficiency and deficiency in Australia (often defined as  $\leq 50\text{nmol/L}$  and  $\leq 38\text{nmol/L}$  25 hydroxyvitamin D<sub>3</sub> respectively). There has been a lack of data from subtropical regions such as Queensland, where there is more intense ultraviolet radiation. Our group has an interest in vitamin D<sup>5</sup>, and had the opportunity to examine vitamin D levels in a case-control study designed to examine risk factors for psychosis.

Within an area of south-east Queensland (latitude 27° south), we randomly sampled individuals with psychosis from a wide range of sites. A well control group was drawn from the same catchment area. Ultraviolet radiation levels were measured using a permanently mounted outdoor erythematous UV meter and sunshine duration was measured with a Campbell-Stokes recorder. Serum 25 hydroxyvitamin D<sub>3</sub> levels were measured with a commercially available radio-immunoassay (DiaSorin Inc). All subjects gave written, informed consent, and the study was approved by the local Institutional Ethics Committee.


The sample consisted of 222 males and 192 females. The mean age was 42.0 years (SD 13.2). 113 subjects lived in “rural areas” as defined by the Australian Bureau of Statistics. Overall the mean (and standard deviation; range) for 25 hydroxyvitamin D<sub>3</sub> in this sample was 69.1 nmol/L (26.2; 12.2-174.5). 23.4% of the sample had levels equal or less than 50 nmol/L, and 8% of the sample had levels equal or less than 38 nmol/L. There was no significant group difference in vitamin D levels between the subjects with psychosis and the well controls ( $t=0.026$ ,  $df=412$ ,  $p = 0.98$ ). Males had significantly higher 25 hydroxyvitamin D<sub>3</sub> levels compared to females (mean, SD; 72.0, 25.8; 65.9, 26.4 nmol/L respectively:  $t=2.36$ ,  $df =412$ ,  $p=0.02$ ). Those living in rural areas had significantly higher 25 hydroxyvitamin D<sub>3</sub> levels compared to those living in urban areas (mean, SD: 74.0, 31.6; 68.1, 23.5 nmol/L respectively:  $t = 2.00$ ,  $df=371$ ,  $p = 0.04$ ).

There was no significant correlation between age and 25 hydroxyvitamin D<sub>3</sub> level (Pearson  $r = -0.06$ ,  $p = 0.25$ ). There were statistically significant positive correlations between 25 hydroxyvitamin D<sub>3</sub> and both sunshine ( $r = 0.10$ ,  $p = 0.04$ ) and ultraviolet radiation ( $r = 0.24$ ,  $p = 0.001$ ). When examined by quartiles of sunshine duration (1<sup>st</sup> quartile being the shortest and 4<sup>th</sup> quartile being the longest duration of sunshine), the mean (SD) 25 hydroxyvitamin D<sub>3</sub> levels for the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> quartiles were not strictly linear - 64.3 (24.9), 66.1 (23.1), 76.3 (24.8) and 67.0 (31.5) respectively.

The within-month variation shown in the Figure was least for July, which also had the lowest mean 25 hydroxyvitamin D<sub>3</sub> levels. We speculate that behaviour during this time of year was modified (less outdoor activity, less skin exposures etc) and/or the weaker ultraviolet radiation during this month was less efficient in producing vitamin D. Living in a subtropical climate does not guarantee adequate vitamin D and supplementation with vitamin D may be more widely needed than previously thought in Queensland.

## References

1. Vasikaran SD, Sturdy G, Musk AA, Flicker L. Vitamin D insufficiency and hyperparathyroidism in Perth blood donors. *Med J Australia* 2000 ;**172**: 406-7.
2. Jones G, Blizzard CL, Riley MD, Paramaeswaran V, Greenway TM, Dwyer T. Vitamin D levels in prepubertal children in Southern Tasmania: prevalence and determinants. *Eur J Clin Nutr* 1999 ;**52**: 824-9.
3. Marks R, Foley PA, Jolley D, Knight KR, Harrison J, Thompson SC. The effect of regular sunscreen use on vitamin D levels in an Australian population. Results of a randomized controlled trial [see comments]. *Arch.Dermatol.* 1995 ;**131**: 415-21.
4. Need AG, Morris HA, Horowitz M, Nordin BEC. Effects of skin thickness, age, body fat, and sunlight on serum 25-hydroxyvitamin D. *Amer J Clin Nutr* 1993 ;**58**: 882-5.
5. McGrath J. Hypothesis: is low prenatal vitamin D a risk-modifying factor for schizophrenia? *Schizophr.Res.* 1999 ;**40**: 173-7.



**Figure 1**

(a) Upper panel. Mean monthly erythemal ultraviolet radiation (joules per square metre) and mean monthly duration of sunshine (hours).

(b) Lower panel. Boxplot of 25 hydroxyvitamin D levels per month (nmol/L). The centre line in the box is the mean, while the upper and low borders of the box define the 25<sup>th</sup> and 75<sup>th</sup> percentile. The whiskers indicate minimum and maximum values excluding outliers, asterisks indicate outliers.